

DOI: <https://doi.org/10.5281/zenodo.14564070>

УДК:616-006:617+616.31-002

OCCURRENCE OF PERIODONTITIS IN WOMEN WITH BREAST CANCER

Nurova Shoxsanam Norpulatovna

Olimov Siddiq Sharifovich

Bukhara State Medical Institute,
Republic of Uzbekistan, Buxara

Abstract: *in women with breast cancer, in women with early menopause, there is a development of a high level of dental diseases in the adaptation phase of the body, with a lack of the hormone estrogen or its abrupt complete absence after radiation, chemotherapy and hormone therapy. It has also been proven that estrogen deficiency negatively affects the condition of the hard tissues of the tooth and especially the condition of periodontal tissues.*

Key words: *dental defects, periodontal disease, hormones, menopause.*

ВОЗНИКНОВЕНИЕ ПАРОДОНТИТА У ЖЕНЩИН С РАКОМ МОЛОЧНОЙ ЖЕЛЕЗЫ

Нурова Шохсанам Норпулатовна

Олимов Сиддик Шарифович

Бухарский государственный медицинский институт
Республика Узбекистан, Бухара,

Резюме: *у женщин с раком молочной железы, у женщин с ранней менопаузой наблюдается развитие высокого уровня стоматологических заболеваний в фазе адаптации организма, при недостатке гормона эстрогена или его резком полном отсутствии после проведенной лучевой, химиотерапии и гормональной терапии. Также было доказано, что дефицит эстрогена негативно влияет на состояние твердых тканей зуба и особенно на состояние тканей пародонта.*

Ключевые слова: *зубочелюстные дефекты, пародонтоз, гормоны, менопауза.*

Menopause is the main cause of chronic generalized periodontitis after radiation and chemical therapy in women with breast cancer. One of the factors of the development of the disease during menopause is changes in the microcirculation system. Microcirculation disorders, chronic hypoxia in periodontal tissues, and nutritional disorders contribute to the destruction of periodontal tissues. On the other hand, microcirculation is primarily caused by inflammation of the tissues of their endothelial walls, ischemia and excessive damage in others, which eliminates the protective barrier, helping the periodontium to become one of the pathomechanisms of the disease.

With long-term treatment, patients with breast cancer develop large and small pain syndromes of the spine of the bones, prolonged and persistent joint pain were observed during dispensary control. The pathogenesis of pain syndrome in patients is often caused by osteoporosis (OP) in breast cancer. Chemotherapy and hormone therapy are factors with a pronounced effect on the metabolic process of the bone structure. [1,3,5].

Chemotherapy also affects bone mineral density by affecting the dysfunction of the ovaries with bone tissue against the background of cytostatic therapy, which at one time causes the rapid development of early menopause and osteoporosis.

In breast cancer (BC), drugs that act on tumors that occur in a hormone-dependent manner affect the patient's menstrual cycle, and at the same time, there is a change in bone density. For adjuvant treatment of breast cancer, tamoxifen hormone antagonists or gonadotropin-releasing antagonists are used, leading to bone loss of up to 7% during one year of treatment [9].

Surgical intervention also leads to artificial menopause according to the indications of ovariectomy, which increases the risk of fractures characteristic of osteoporosis. [8]. Thus, changes in bones caused by the use of endocrine therapy, chemotherapy and surgical methods in the treatment of breast cancer are currently an urgent problem. [9]. Osteoporosis causes unpleasant symptoms, negatively affecting patients with this type of cancer.[6]. Solving the problem of the spread of periodontal tissue diseases is one of the most important and urgent tasks of modern dentistry. 12% of the population has a healthy periodontal disease. In people over 35 years of age, the proportion of initial periodontal changes gradually decreases by 26-15%, at the same time, an increase in the number of changes from moderate to severe to 75% was observed. This, in turn, leads to the appearance of functional disorders in the dental and maxillary system. According to WHO, tooth loss as a result of periodontal diseases, functional disorders of the dental system develop five times faster than complications of caries, and is the second most common among all dental diseases. According to WHO, the incidence of periodontal disease is 55-69% in adolescents, 86.2% at the age

of 35-44 years, and increases significantly at the age of 40-50 years (65-98%). In this regard, the diagnosis, treatment and prevention of these diseases will be of general medical and social importance. The issues of etiology and pathogenesis of periodontal diseases are explained in terms of the general influence of local and widespread factors, among them hormonal changes are pronounced, especially during postmenopause. It has been proven that insufficient estrogen levels during postmenopause cause an imbalance between resorption and bone formation, and this leads to rapid changes in bone structure, the development of osteoporosis and its complications. At the present stage, the relationship between the bone mineral density in various parts of the bones of the body and the height of the alveoli in the interproximal sections has been determined. According to the results of the study, a decrease in bone mineral density leads to a decrease in the interalveolar height, to a bend in the alveolar process that attaches to the gum or to the transitional fold. Osteoporosis during postmenopause is a dangerous factor in the development of periodontal diseases.

The study was conducted in 2021-2023 at the Bukhara branch of the Republican Specialized Scientific and Practical Medical Center of Oncology and Radiology and at the Dental Scientific and Practical Center at the Bukhara State Medical Institute. The total number of examined patients was 703, of which 154 were control patients. with

154 patients aged 55 to 65 years with diffuse periodontitis of moderate and severe severity, whose postmenopause lasted at least 5 years, were examined and treated. The examined patients are divided into groups:

Group 1 consisted of 55 patients with CKD on the background of general osteoporosis (average age 58.30 ± 0.65), menopause age 49.23 ± 0.85 , menopause duration was noted.

Group 2 of 40 patients with CKD on the background of osteopenia (mean age 57.73 ± 0.42 years), menopause age 48.65 ± 0.69 , menopause duration 9.5 ± 0.88 .

In the comparative group, bone mineral density was normal (average age 57.92 ± 0.49 years) at menopausal age 48.37 ± 0.92 years, with menopause duration 9.47 ± 0.59 years, in 40 patients with moderate menopausal CKD, and with moderate and severe CKD aged 25 to 45 years, 30 patients are in reproductive age (average age 39.5 ± 1.2 years), in whom bone tissue disorders were not noted. 20 women of reproductive age from the control group are practically healthy (average age 37.8 ± 1.1).

To participate in the study, patients who had been diagnosed with breast cancer for at least 5 years, postmenopausal women with osteoporosis or CKD on the background of osteopenic syndrome were involved, and consent was obtained from them to participate in the study.

Changes in the oral cavity in HDPE in patients with breast cancer

Indicators	Group of patients	
	Women with perimenopausal breast cancer n=57	Postmenopausal women with breast cancer n= 39
Hygiene Index (OHI-S)	<u>2,27±0,05</u> 2,18±0,05	<u>2,55±0,07*</u> 2,32±0,04*
Degree of bleeding	<u>2,29±0,07</u> 2,52±0,08#	<u>2,45±0,04*</u> 2,88±0,06*#
RMA	<u>50,32±1,79</u> 55,43±2,15#	<u>61,68±1,93*</u> 69,88±2,17*#
PI	<u>4,82±0,15</u> 4,56±0,18#	<u>6,12±0,13*</u> 5,87±0,15*#
The depth of the periodontal pocket, mm	<u>5,22±0,21</u> 4,75±0,15	<u>6,75±0,44*</u> 5,49±0,27*
Fuchs index for the upper jaw	<u>0,58±0,06</u> 0,60±0,03	<u>0,45±0,04*</u> 0,48±0,03*
Fuchs index for the lower jaw	<u>0,60±0,04</u> 0,62±0,03	<u>0,48±0,03*</u> 0,50±0,03*
Cortical index	<u>4,68±0,03</u> 4,70±0,05	<u>4,40±0,03*</u> 4,55±0,04*

The objective condition of periodontal tissues was studied based on the dynamics of the following tests:

- simplified hygienic index (OHI-S);
- determination of the level of bleeding gums (Cowell I., 1975).
- - papillary-marginal-alveolar index (PMA) (Parma G., 1960);
- periodontal index (Russel A., 1967);
- measurement of the depth of the periodontal pocket – (WHO data, 1989)
- determination of the degree of pathological tooth movement (Fleszar T. J. et al., 1980).

A clinical examination of the oral cavity of women with breast cancer was conducted in order to determine the condition of the teeth, hard tissues of teeth, type of bite, protruding fold, presence of subgingival and supra- gingival solid carious deposits, condition of the oral mucosa (edema, hyperemia, bleeding), exudate condition, presence and depth of periodontal pockets, determination of the degree of pathological mobility of teeth. Attention was paid to the timing of the

appearance of inflammatory changes in periodontal tissue and their connection with the onset of menopause. The results of the survey are presented in the table below.

A clinical examination of the oral cavity of women with breast cancer was conducted in order to determine the condition of the teeth, hard tissues of teeth, type of bite, protruding fold, presence of subgingival and supra-gingival solid carious deposits, condition of the oral mucosa (edema, hyperemia, bleeding), exudate condition, presence and depth of periodontal pockets, determination of the degree of pathological mobility of teeth. Special attention was paid to the timing of the appearance of inflammatory changes in periodontal tissues and their connection with the onset of menopause. The results of the survey are presented in the table below. Расчет индекса Фукса осуществляется по следующей формуле:

$$((n \times 0) + (pc \ 1) + (pc \ 2) + (pc \ 3) + (pc \ 4) \text{ number of teeth})$$

Index of characteristics of the cortical layer of the mandible Klemetti E. Et al. was carried out on the basis of the MCI index. It was evaluated according to the morphological characteristics of the bone. Depending on its morphological characteristics, 3 types were distinguished: C 1 - the inner boundary of the normal cortical layer is clear, the cortical region is unknown without changes. C 2 - one side of the cortical plate has the shape of a crescent and is significantly damaged cortically. C 3-the cortical edge is uneven, the cortical plate is multilayered with a large number of defective areas. The initial orthopantomograms were also analyzed 2 years after treatment.

Chronic diffuse periodontitis in patients with breast cancer (%)

Characteristics of damaged periodontal tissue in patients with breast cancer	Group of patients		
	Healthy n=40(%)	osteopenia, n=40(%)	Osteoporosis n=60(%)
Chronic diffuse periodontitis of medium form	23 (57,5)	26 (65,0)	36 (60,0)
Severe chronic diffuse periodontitis	17 (42,5)	14 (35,0)	24 (40,0)

Among patients with chronic diffuse periodontitis and postmenopausal osteoporosis, according to clinical and radiological data, 60% had moderate periodontal disease, and 40% had severe periodontal disease. Among the examined patients, with an average periodontal disease of 65%, severe periodontitis was diagnosed in 35% against the background of osteopenia. No significant difference was found in the statistical comparison of patients by group.

LITERATURES

1. Nurova Sh.N. Treatment and Prevention of Dental Deformities in Women in Early Menopause, MEDICAL AND NATURAL SCIENCES Volume: 03 Issue: 03 | May- Jun 2022 ISSN: 2660-4159
2. Нурова Ш.Н. Обзорный характеристика об этиологии, диагностики, лечение и профилактики зубочелюстных деформаций у женщин в ранней менопаузы. Oral medicine and craniofacial research Volume: 03 Issue: 1| 2022 ISSN: 2181-096623-26
3. Nurova Sh.N. Prediction and Prevention of the Development of Partial and Complete Defects of the Dentition in Women in Early Menopause. MIDDLE EUROPEAN SCIENTIFIC BULLETIN Volume: 28, ISSN 2694-9970 11-15
4. Нурова Ш.Н. Аёлларда эстроген етишмовчилиги оқибатида тиш-жағ тизимидаги ўзгаришлар бўйича умумий тавсиф. Oral medicine and craniofacial research Volume: 4 Issue: 2| 2023 ISSN: 2181-0966. Б. 53-56
5. Nurova Sh.N. General Description of Changes in the Gums in Women with Breast Cancer / RESEARCH JOURNAL OF TRAUMA AND DISABILITY STUDIES Volume: 2, ISSN 2720-6866 220-224
6. Scardina, G.A. Oral microcirculation in post-menopause: a possible correlation with periodontitis / G.A. Scardina, P. Messina // Gerodontology. - 2012. - Vol.29(2). - P.e1045-1051.
6. Ramya, K.S. Expression of VEGF in Periodontal Tissues of Type II Diabetes Mellitus Patients with Chronic Periodontitis -an Immunohistochemical Study / K.S. Ramya // J Clin Diagn Res. - 2014. - Vol.8(8). - P. ZC01-3.
7. Нуров Н.Б. Analysis of data on the morphological structure and biomechanics of the temporomandibularis system World Bulletin of Public Health (WBPH) ISSN 2749– 3644 Volume 3, October-2021; 85-87 Impact factor: 7.635
8. Нуров Н.Б. Prevalence of adentia in the elderly and development factors. World Bulletin of Public Health (WBPH) ISSN 2749– 3644 Volume 3, October-2021; 78-81 Impact factor: 7.635
9. Нуров Н.Б. Treatment of old people according to age specialties World Bulletin of Public Health (WBPH) ISSN 2749– 361X Volume 3, October-2021; 125-128 Impact factor: 7.545
10. Нуров Н.Б., Тешаев Ш.Ж., Олимов С.Ш. Morphological features of the face-jaw in humans with complete and partial adentia Journal of Advanced Zoology ISSN: 0253 7214 Volume 44, Issue S-2 Year 2023 Page 770:778